

# GLOBAL WELL-POSEDNESS IN SOBOLEV SPACE IMPLIES GLOBAL EXISTENCE FOR WEIGHTED $L^2$ INITIAL DATA FOR $L^2$ -CRITICAL NLS.

P. BLUE, J. COLLIANDER

**ABSTRACT.** The  $L^2$ -critical defocusing nonlinear Schrödinger initial value problem on  $\mathbb{R}^d$  is known to be locally well-posed for initial data in  $L^2$ . Hamiltonian conservation and the pseudoconformal transformation show that global well-posedness holds for initial data  $u_0$  in Sobolev  $H^1$  and for data in the weighted space  $(1+|x|)u_0 \in L^2$ . For the  $d = 2$  problem, it is known that global existence holds for data in  $H^s$  and also for data in the weighted space  $(1+|x|)^\sigma u_0 \in L^2$  for certain  $s, \sigma < 1$ . We prove: If global well-posedness holds in  $H^s$  then global existence and scattering holds for initial data in the weighted space with  $\sigma = s$ .

## 1. INTRODUCTION

Consider the initial value problem for the  $L^2$  critical nonlinear Schrödinger equation for  $u : \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{C}$ ,

$$(1.1) \quad \begin{cases} i\partial_t u + \Delta u = \lambda |u|^{\frac{4}{d}} u \\ u(t_0, x) = u_0(x). \end{cases}$$

This problem is called *defocusing* for  $\lambda > 0$  and *focusing* for  $\lambda < 0$ . In dimension  $d = 2$ , equation (1.1) reduces to the cubic nonlinear Schrödinger equation,  $i\partial_t u + \Delta u = \lambda |u|^2 u$ , which appears widely as a model equation in Physics [9].

This problem is locally well-posed in  $L^2$  or in any  $H^s$  with  $s \geq 0$ . That is, given initial data  $u_0 \in H^s(\mathbb{R}^d)$  with  $s \geq 0$  there is a local existence time,  $T_{lwp}$ , and a local in time solution  $u : [t_0 - T_{lwp}, t_0 + T_{lwp}] \times \mathbb{R}^d \rightarrow \mathbb{C}$  such that  $u$  solves (1.1) and the function  $u \in C_t H^s$ . For  $s > 0$ ,  $T_{lwp}$  is a decreasing function of the  $H^s$  norm of  $u_0$ .

An open problem is to prove global well-posedness in  $L^2$  in the defocusing case and under an appropriate smallness condition in the focusing case. For the focusing case, it is believed that solutions with  $L^2$  norm smaller than the ground state<sup>1</sup> mass  $\|Q\|_{L^2}$  do not blowup and in fact scatter. Explicit blow-up solutions with Schwartz class initial data and with the mass of the ground state are known to exist in the focusing case. By finding the optimal constant in the Gagliardo-Nirenberg estimate,  $\|u\|_{L^{\frac{2(d+2)}{d}}}^{\frac{2(d+2)}{d}} \leq (\frac{d+2}{d} \|Q\|_{L^2}^{-\frac{4}{d}}) \|\nabla u\|_{L^2}^2 \|u\|_{L^2}^{\frac{4}{d}}$ , Weinstein [11] proved that  $H^1$  initial data with  $L^2$  norm less than the ground state mass evolves globally in time. In the defocusing case,  $L^2$  solutions are expected to exist globally in time and scatter. Although the  $L^2$  norm of  $u(t)$  is constant on the local well-posedness time interval, this norm does not control the length of the local well-posedness time  $T_{lwp}$  and

---

Date: January 9, 2010.

<sup>1</sup>The *ground state* is the unique (up to translations) positive solution of  $-Q + \Delta Q = Q^3$ .

can not be used to prove global well-posedness in  $L^2$ . In  $H^1$ , this problem has an additional conserved quantity, the energy,  $E[u] = \int \frac{1}{2} |\nabla u|^2 + \frac{\lambda}{4} |u|^4 dx$ . In the defocusing case, the energy is positive and dominates the  $H^1$  norm. Since the local well-posedness time is a function of the  $H^1$  norm, at any time the solution persists for a uniformly long local well-posedness time, and, hence, globally in time.

Bourgain was the first to prove [2], for the cubic problem in  $\mathbb{R}^2$ , global well-posedness below the energy threshold  $H^1$  by proving global well-posedness for data in  $H^s$  in the defocusing case for  $s > \frac{3}{5}$ . The method in [2] involved a decomposition of the data into high and low frequencies with a sharp cut-off function in the Fourier variables. Later, the “ $I$ -method” [6] was used to improve this to  $s > \frac{4}{7}$  for the cubic problem on  $\mathbb{R}^2$ .

The nonlinear Schrödinger equation (1.1) has a discrete pseudoconformal symmetry,

$$(1.2) \quad \mathcal{C}[u](\tau, y) = v(\tau, y) = \frac{1}{|\tau|^{\frac{d}{2}}} e^{\frac{iy^2}{4\tau}} u\left(-\frac{1}{\tau}, \frac{y}{\tau}\right).$$

This is a symmetry in the sense that, if  $u(t, x)$  is a solution to the nonlinear Schrödinger equation on  $(t, x) \in [t_1, t_2] \times \mathbb{R}^d$ , then  $v(\tau, y)$  is a solution on  $\tau \in [-t_1^{-1}, -t_2^{-1}]$ ,  $y \in \mathbb{R}^d$ . Throughout this paper, we shall use  $u$ ,  $t$ , and  $x$  to refer to a solution, the time variable, and the spatial variable respectively, to use  $v$  as the pseudoconformal transform of  $u$ , and to use  $\tau$  and  $y$  as the arguments of  $v$  which will be called the transformed time and space variables. In particular,  $\tau = t^{-1}$ .

We introduce the space

$$(1.3) \quad H^{0,s} = \{u \in L^2 : |x|^s u \in L^2\}$$

with norm

$$(1.4) \quad \|u\|_{H^{0,s}} = \|\langle x \rangle^s u\|_{L^2}$$

where  $\langle x \rangle^2 = 1 + x^2$ .

The pseudoconformal transform satisfies

$$(1.5) \quad E[v](\tau) = \|xu_0\|_{L^2}^2 = \|u_0\|_{H^{0,1}}^2.$$

The energy on the left side of this equation is already known to be independent of  $\tau$ . This property was used to prove global well-posedness in  $L^2$  for initial data  $u_0 \in H^{0,s}$  with  $s > \frac{3}{5}$  in the defocusing case [2]. The proof involves a spatial decomposition analogous to the Fourier decomposition used in proving the  $H^s$  global existence result. As a further consequence, [2] also proves scattering, that is the existence of functions  $u_{\pm} \in L^2$  such that

$$\lim_{t \rightarrow \pm\infty} \|u(t) - e^{\pm it\Delta} u_{\pm}\|_{L^2} = 0.$$

This paper establishes that global well-posedness in  $H^s$  for (1.1) implies global existence and scattering in  $L^2$  for initial data in  $H^{0,s}$  for (1.1). Thus, the link between  $H^s$  global well-posedness and the evolution properties of  $H^{0,s}$  initial data found in the  $\mathbb{R}^2$  case in [2] is in fact common to all pseudoconformal or  $L^2$ -critical nonlinear Schrödinger initial value problems (1.1).

**Proposition 1.1.** *Assume that the nonlinear Schrödinger equation (1.1) is globally well-posed in  $H^s$  (with the additional hypothesis that the initial data has  $L^2$  norm bounded by  $\|Q\|_{L^2}$  in the focusing case).*

If  $u_0 \in H^{0,s}$  (and  $\|u_0\|_{L^2} < \|Q\|_{L^2}$  in the focusing case), then there is a function  $u : \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{C}$  which solves the nonlinear Schrödinger equation (1.1) for all time. Furthermore, there are functions  $u_{\pm} \in H^{0,s}$  such that

$$\lim_{t \rightarrow \pm\infty} \|e^{\mp it\Delta} u(t) - u_{\pm}\|_{H^{0,s}} = 0.$$

Based on the  $H^s$  global well-posedness result in [6], we obtain as a consequence of Proposition 1.1 that for  $s > \frac{4}{7}$ , initial data in  $H^{0,s}$  evolve globally in time and scatter in  $L^2$  under the cubic nonlinear Schrödinger flow on  $\mathbb{R}^2$ .

Global existence in  $L^2$  for initial data in  $H^{0,s}$  means initial data  $u_0 \in H^{0,s}$ , which is also in  $L^2$ , evolves as a solution in  $L^2$  and that this solution exists for all time. This is significantly weaker than global well-posedness in  $H^s$ , which means that initial data in  $H^s$  continuously evolves in  $H^s$ , that this solution exists for all time, and that the time evolution map  $S_{NLS}(t, 0) : u_0 \mapsto u(t)$  is continuous from  $H^s$  to  $H^s$ .

The asymmetry between  $H^s$  and  $H^{0,s}$  in Proposition 1.1 is a consequence of the local theory. The initial value problem (1.1) is locally well-posed in  $H^s$ ; whereas, in  $H^{0,s}$ , (1.1) is ill-posed, so global existence in  $L^2$  for initial data in  $H^{0,s}$  can not be extended to  $H^{0,s}$  global well-posedness.

In the remainder of the introduction, we review  $L^2$  and  $H^s$  local-well posedness,  $H^{0,s}$  ill-posedness, and some properties of the pseudoconformal transform. In Section 2, we prove Proposition 1.1 by showing that, for a solution with initial data in  $H^{0,s}$ , the pseudoconformal transform is in  $H^s$ . This is done by taking regularized approximators and showing their transforms converge in  $H^s$  at a particular transformed time  $-T_{lwp}^{-1}$ . In Section 3, we show that scattering is a consequence of the construction in Section 2.

We use the notation  $S_{NLS}(t_2, t_1)$  to denote the nonlinear Schrödinger evolution map from time  $t_1$  to time  $t_2$ ,  $\mathcal{F}[\bullet]$  for the Fourier transform, and  $\Re\alpha$  and  $\Im\alpha$  for the real and imaginary part of  $\alpha$  respectively.

**1.1. Local well-posedness theory.** The local well-posedness theory (see [3], [8] for a review) begins with the presentation of the nonlinear Schrödinger equation as an integral equation through Duhamel's principle

$$(1.6) \quad u = \Phi_{u_0}[u],$$

$$(1.7) \quad \Phi_{u_0}[u] = e^{it\Delta} u_0 + \int_0^t e^{i(t-t')\Delta} (\lambda |u|^2 u)(t') dt'.$$

To prove that (1.6) has a unique solution, it is sufficient to show that  $\Phi_{u_0}$  is a contraction in an appropriate space. This space will be the Strichartz space defined below.

**Definition 1.2.** The pair  $(q, r)$  is  $L^2$  Strichartz admissible, or simply admissible, if

$$\frac{2}{q} + \frac{d}{r} = \frac{d}{2}, \quad 2 \leq q \leq \infty, \quad 2 \leq r \leq 2 + \frac{4}{d-2}.$$

In dimension  $d = 2$ , there is the additional restriction that  $2 < q \leq \infty$  and  $2 \leq r < \infty$ . In dimension  $d = 1$ ,  $2 \leq q \leq \infty$  and  $2 \leq r \leq \infty$ .

For  $s \geq 0$ , a pair  $(q, r)$  is  $H^s$  Strichartz admissible if

$$\frac{2}{q} + \frac{d}{r} = \frac{d}{2} - s, \quad 2 \leq q \leq \infty, \quad 2 \leq r.$$

The  $L^2$  Strichartz norm, or simply the Strichartz norm, is

$$\|u\|_{S^0} = \sup_{(q,r) \text{ admissible}} \|u\|_{L_t^q L_x^r}.$$

The homogeneous  $H^s$  Strichartz norm is

$$\|u\|_{\dot{S}^s} = \sup_{(q,r) \text{ admissible}} \|D^s u\|_{L_t^q L_x^r},$$

where  $D$  is the Fourier multiplier defined by  $\mathcal{F}[Df](\xi) = |\xi| \mathcal{F}[f](\xi)$ .

For  $s > 0$ , the  $H^s$  Strichartz norm is

$$\|u\|_{S^s} = \sup_{0 \leq \sigma \leq s} \|u\|_{\dot{S}^\sigma}$$

For an interval  $I$ , the spaces  $S^0(I)$ ,  $\dot{S}^s(I)$ , and  $S^s(I)$  are the spaces with the above Strichartz norm where the  $t$  integration is taken over the interval  $t \in I$ .

With this notation, we record the Strichartz estimates: For  $s > 0$ ,

$$\begin{aligned} \|e^{it\Delta} u_0\|_{S^0} &\leq C \|u_0\|_{L^2}, \\ \|e^{it\Delta} u_0\|_{\dot{S}^s} &\leq C \|u_0\|_{\dot{H}^s}, \\ \|e^{it\Delta} u_0\|_{S^s} &\leq C \|u_0\|_{H^s}. \end{aligned}$$

Using Duhamel's principle, estimates for solutions to the linear Schrödinger equation, and a contraction argument, it is possible to prove local well-posedness, uniqueness, and continuity of solutions in the  $L^2$  Strichartz space. The balanced pair  $q = r = \frac{2(d+2)}{d}$  plays an important role in our presentation of the local well-posedness theory.

**Theorem 1.3** ( $L^2$  local well-posedness). *For  $u_0 \in L^2$  and  $\delta_1 > 0$ , there is a  $T_{lwp} = T_{lwp}(u_0, \delta_1) > 0$  and  $u : [0, T_{lwp}] \times \mathbb{R}^d \rightarrow \mathbb{C}$  such that:*

- (1)  $u$  solves the nonlinear Schrödinger equation (1.1) with initial data  $u_0 \forall t \in [0, T_{lwp}]$ ;
- (2)  $\|u\|_{S_{([0, T_{lwp}])}^0} \leq 2\|u_0\|_{L^2}$ ;
- (3)  $\|u\|_{L_{tx}^{\frac{2(d+2)}{d}}([0, T_{lwp}])} \leq \delta_1$ ;
- (4) for all  $t \in [0, T_{lwp}]$ ,  $\|u(t)\|_{L^2} = \|u_0\|_{L^2}$ ;
- (5) there is a  $\delta_2 > 0$ , such that, if  $u'_0 \in L^2$  with  $\|u_0 - u'_0\|_{L^2} < \delta_2$ , then there is a  $u' \in S^0([0, T_{lwp}])$  such that  $u'$  solves the nonlinear Schrödinger equation with initial data  $u'_0$ ; and  $\|u - u'\|_{S^0([0, T_{lwp}])} \leq 2\|u_0 - u'_0\|_{L^2}$ . In particular, for  $t \in [0, T_{lwp}]$ ,

$$(1.8) \quad \|u(t) - u'(t)\|_{L^2} \leq 2\|u_0 - u'_0\|_{L^2}.$$

The  $T_{lwp}$  from this theorem will be referred to as the  $L^2$  local well-posedness time. The  $L^2$  maximal forward time of existence,  $T^*$ , is the time for which there is a solution  $u : [0, T^*) \rightarrow L^2$  but no solution  $\tilde{u} : [0, t] \rightarrow L^2$  for  $t > T^*$ .

The  $L^2$  local well-posedness theory and  $L_{tx}^{\frac{2(d+2)}{d}}([0, T_{lwp}])$  norm play a central role in our presentation of the  $H^s$  theory. The  $L^2$  local well-posedness time,  $T_{lwp}$ , is the

same as the  $H^s$  local well-posedness time. If a solution has small  $L_{tx}^{\frac{2(d+2)}{d}}([0, T_{lwp}])$  norm, then its  $H^s$  norm can not grow by more than a small factor, and, if another solution has initial data close in  $L^2$ , then the  $H^s$  distance between the two solutions can not grow by more than a small factor. The  $L^2$  local well-posedness Theorem 1.3 uses  $T_{lwp}$  chosen to enforce the  $L_{tx}^{\frac{2(d+2)}{d}}([0, T_{lwp}])$  smallness condition.

**Theorem 1.4** ( $H^s$  local well-posedness). *For  $u_0 \in H^s$ , the solution  $u$  and local well-posedness time  $T_{lwp}$  given in Theorem 1.3 satisfy:*

- (1) for  $t \in [0, T_{lwp}]$ ,  $u(t) \in H^s$ ;
- (2) there is a  $\delta_3 > 0$ , such that, if  $\|u\|_{L_{tx}^{\frac{2(d+2)}{d}}(I)} \leq \delta_3$ , then  $\|u\|_{\dot{S}^s(I)} \leq 2\|u_0\|_{H^s}$ ;
- and
- (3) there is a  $\delta_4 > 0$ , such that, if  $\|u\|_{L_{tx}^{\frac{2(d+2)}{d}}(I)} \leq \delta_3$ ,  $u'$  is a solution to the nonlinear Schrödinger equation with initial data  $u'_0 \in H^s$ , and  $\|u_0 - u'_0\|_{L^2} \leq \delta_4$ , then  $\|u - u'\|_{S^s} \leq 2\|u_0 - u'_0\|_{H^s}$ . In particular,

$$(1.9) \quad \|u(t) - u'(t)\|_{H^s} \leq 2\|u_0 - u'_0\|_{H^s}.$$

Since  $u$  can be extended to any interval on which the  $L^2$  Strichartz norm is finite, the  $L^2$  Strichartz norm must diverge on intervals approaching the maximal forward time of existence.

**Corollary 1.5** (Maximal time blow up theorem). *If  $u$  is a solution to the nonlinear Schrödinger equation (1.1) with initial data  $u_0 \in L^2$  and maximal forward time of existence  $T^*$  (that is  $u : [0, T^*) \rightarrow \mathbb{C}$  is a solution, but for  $t > T^*$ , there is no solution  $\tilde{u} : [0, t] \rightarrow \mathbb{C}$ ), then*

$$(1.10) \quad \lim_{t \uparrow T^*} \|u\|_{S^0([0, t])} = \infty.$$

**1.2. Local evolution of  $H^{0,s}$  initial data.** As noted earlier, Proposition 1.1 only asserts global existence in  $L^2$  for initial data in  $H^{0,s}$ , not  $H^{0,s}$  global well-posedness, because the nonlinear Schrödinger equation is ill-posed in  $H^{0,s}$ . This ill-posedness is a consequence of the ill-posedness of the linear Schrödinger equation in  $H^{0,s}$ . It has solutions which start with finite initial  $H^{0,s}$  norm, but have divergent  $H^{0,s}$  norm at all later times. From these, it is possible to show that the nonlinear Schrödinger time evolution  $S_{NLS}(t, 0) : u_0 \mapsto u(t)$  is not continuous from  $H^{0,s}$  to  $H^{0,s}$ . This result does not have an analogue in  $H^s$ ,  $s > 0$ .

Solutions to the linear Schrödinger equation which have finite initial  $H^{0,s}$  norm but divergent  $H^{0,s}$  norm can be constructed from sums of Gaussians. Since the Fourier variable corresponds to velocity for the linear Schrödinger equation, Gaussian initial data of width  $a^{-\frac{1}{2}}$  has a Gaussian Fourier transform with a width  $a^{\frac{1}{2}}$ , and hence disperses linearly in time at speed  $a^{\frac{1}{2}}$ . A linear combination of such initial data can be constructed to disperse with unbounded velocity and instantaneously divergent weighted norm.

**Lemma 1.6.** *For  $s > 0$ , there is a solution  $\Psi$  to the linear Schrödinger equation with initial data  $\Psi_0 \in H^{0,s}$  for which  $\Psi(t)$  is not in  $H^{0,s}$  at any future time  $t > 0$ .*

*Proof.* Let

$$\begin{aligned} (C_{d,s})^2 &= \frac{1}{\pi^{\frac{d}{2}}} \int_{\mathbb{R}^d} |x|^{2s} e^{-x^2} dx, \\ (C'_{d,s})^2 &= \frac{1}{\pi^{\frac{d}{2}}} \int_{|x|<1} |x|^{2s} e^{-x^2} dx, \\ r(a,t) &= \left( \frac{1+4a^2t^2}{a} \right)^{\frac{1}{2}}, \end{aligned}$$

and let  $\chi_{|x|<r(a,t)}$  be the characteristic function with support on  $|x| < r(a,t)$ .

Given  $A > 0$  and  $a > 0$ , let  $\psi[A, a]$  be solutions to the linear Schrödinger with initial data

$$u_0[A, a] = A \left( \frac{a}{\pi} \right)^{\frac{d}{4}} e^{\frac{-ax^2}{2}}.$$

These are given by

$$\begin{aligned} \psi[A, a] &= e^{it\Delta} u_0[A, a] = A \left( \frac{a}{\pi} \right)^{\frac{d}{4}} \frac{1}{(1+2ait)^{\frac{d}{2}}} e^{\frac{-ax^2}{2(1+2ait)}} \\ \|e^{it\Delta} u_0[A, a]\|_{L_x^2} &= A \\ \| |x|^s e^{it\Delta} u_0[A, a] \|_{L_x^2} &= C_{d,s} A r(a,t)^s \sim A a^{\frac{s}{2}} t^s \quad \text{for large } t \end{aligned}$$

Let  $\psi_k = e^{it\Delta} u_0[A_k, a_k]$  with  $A_j$  chosen, in terms of  $A_k$  for  $k < j$ , to be sufficiently small so that

$$(1.11) \quad \sum_{i=k+1}^j A_i \leq \frac{1}{8} C'_{d,s} A_k$$

and with  $a_j$  chosen sufficiently large so that

$$(1.12) \quad \sum_{k < j} A_k (1 + C_{d,s} r(a_k, t)^s) \leq \frac{1}{4} C'_{d,s} A_j a_j^{\frac{s}{2}} t^s,$$

and let  $\Psi = \sum \psi_k$ .

Given a fixed  $t$ , let  $r_k = r(a_k, t)$  and  $\chi_k = \chi_{|x|<r_k}$ . The first condition, (1.11), ensures that, for sufficiently large  $k$ , on a length scale of  $|x| < r_k$ , the function  $\psi_k$  dominates all the later  $\psi_j$  with  $j > k$ :

$$\|\chi_k |x|^s \psi_k\|_{L^2} = C'_{d,s} A_k r_k^s > 8 \sum_{j>k} A_j r_k^s > 8 r_k^s \sum_{j>k} \|\psi_j\|_{L^2} > 4 \sum_{j>k} \|\chi_k |x|^s \psi_j\|_{L^2}$$

The second condition, (1.12), ensures that  $A_j a_j^{\frac{s}{2}}$  grows at least exponentially. It also ensures that, for  $a_j^{\frac{1}{2}} > t^{-2}$ , in  $H^{0,s}$ ,  $\psi_j$  dominates all of the previous  $\psi_k$  with  $k < j$ :

$$\|\chi_j |x|^2 \psi_j\|_{L^2} = C'_{d,s} A_j r_j^s > C'_{d,s} A_j a_j^{\frac{s}{2}} t^s > 4 \sum_{k<j} A_k (1 + C_{d,s} r_k^s) > 4 \sum_{k<j} \|\psi_k\|_{H^{0,s}}$$

Since  $\Psi$  is the sum of the  $\psi_k$ , and since, at a given time  $t$ , for sufficiently large  $j$ , on a length scale of  $r_j$ ,  $\psi_j$  dominates all the other  $\psi_k$ , the  $H^{0,s}$  norm of  $\Psi(t)$  is bounded below by arbitrarily large numbers and must diverge:

$$\|\Psi(t)\|_{H^{0,s}} \geq \|\chi_j |x|^s \Psi\|_{L^2} \geq \|\chi_j |x|^s \psi_j\|_{L^2} - \sum_{k \neq j} \|\chi_j |x|^s \psi_k\|_{L^2} \geq \frac{1}{2} \|\chi_j |x|^s \psi_j\|_{L^2} \geq C'_{d,s} A_j a_j^{\frac{s}{2}} t^s \rightarrow \infty.$$

□

A similar sequence of Gaussian initial data shows that the nonlinear Schrödinger equation is also ill-posed in  $H^{0,s}$ .

**Proposition 1.7.** *The nonlinear Schrödinger equation (1.1) is not well-posed in  $H^{0,s}$  for any  $s > 0$ : the nonlinear Schrödinger evolution from time 0 to time  $t > 0$ ,  $S_{NLS}(t, 0) : u_0 \mapsto u(t)$ , is not continuous from  $H^{0,s}$  to  $H^{0,s}$ .*

*Proof.* It is sufficient to construct a sequence of  $L^2$  solutions,  $u^{[k]}$ , for which  $\|u_0^{[k]}\|_{H^{0,s}} \rightarrow 0$ , but for which, at any  $t > 0$ ,  $\|u^{[k]}(t)\|_{H^{0,s}} \rightarrow \infty$ . This construction follows from the closeness of nonlinear Schrödinger and linear Schrödinger evolutions for small initial data and from the existence of linear solutions with arbitrarily fast  $H^{0,s}$  growth.

When the  $L_{tx}^{\frac{2(d+2)}{d}}$  norm is sufficiently small, the difference between the linear Schrödinger and nonlinear Schrödinger evolutions is small. From Duhamel's principle and an extension of the local well-posedness theory, it is known that there is a  $\delta'$  such that, if  $\|u_0\|_{L^2} \leq \delta'$ , then  $u$  is defined for all time, and

$$\begin{aligned} \|u - e^{it\Delta}u_0\|_{S^0} &\leq \|u\|_{L_{tx}^{\frac{2(d+2)}{d}}}^{\frac{4}{d}+1}, \\ \|u\|_{L_{tx}^{\frac{2(d+2)}{d}}} &\leq 2\|u_0\|_{L^2}. \end{aligned}$$

For the linear Schrödinger solutions, the notation from the previous lemma will be used. In addition,  $u^{[k]}$  will denote the nonlinear Schrödinger evolution of  $u_0[A_k, a_k]$  with  $A_k$  decreasing to zero and  $a_k$  increasing to infinite, but with rates to be chosen. The index  $k$  will be chosen sufficiently large so that, for  $i > k$ ,

$$\begin{aligned} A_i &< \delta' \\ A_i^{\frac{4}{d}} &< \frac{C'_{d,s}}{2^{\frac{4}{d}+2}} \end{aligned}$$

As in Lemma 1.6, the  $H^{0,s}$  norm can be estimated by localizing on a length scale of  $r_k$ .

$$\begin{aligned} \|u^{[k]}(t)\|_{H^{0,s}} &\geq \|\chi_k |x|^s u^{[k]}\|_{L^2} \geq \|\chi_k |x|^s e^{it\Delta} u_0^{[k]}\|_{L^2} - \|\chi_k |x|^s (u^{[k]} - e^{it\Delta} u_0^{[k]})\|_{L^2} \\ &\geq C'_{d,s} A_k r_k^s - r_k^s 2^{\frac{4}{d}} \|u_0^{[k]}\|_{L^2}^{\frac{4}{d}} \geq C'_{d,s} A_k r_k^s (1 - \frac{1}{2}) \gtrsim A_k a_k^{\frac{s}{2}} t^s. \end{aligned}$$

If  $A_k = a_k^{-\frac{s}{4}}$  and  $a_k \rightarrow \infty$ , then, for each  $t > 0$ ,

$$\begin{aligned} \|u_0^{[k]}\|_{H^{0,s}} &\sim A_k \rightarrow 0, \\ \|u^{[k]}(t)\|_{H^{0,s}} &\sim A_k a_k^{\frac{s}{2}} t^s \rightarrow \infty. \end{aligned}$$

□

**1.3. The pseudoconformal transform and Strichartz norms.** The pseudoconformal transform is a symmetry of both the linear Schrödinger equation and the pseudoconformal nonlinear Schrödinger equation (1.1) and is also an isometry on  $L_x^2$  and the Strichartz admissible  $L^q L^r$  spaces.

**Lemma 1.8.** (1) *If  $0 \leq t_1 \leq t_2$ , and  $u : [t_1, t_2] \times \mathbb{R}^d \rightarrow \mathbb{C}$  satisfies the nonlinear Schrödinger equation (1.1), then so does  $\mathcal{C}[u] = v : [-t_1^{-1}, -t_2^{-1}] \times \mathbb{R}^d \rightarrow \mathbb{C}$ .*




FIGURE 1. The map  $F_t : u_0 \mapsto v(-T_{lwp}^{-1})$ .  $u_0$  is given in  $H^{0,s} \subset L^2$ , and  $u'_0$  and  $u''_0$  in  $H^{0,s} \cap H^1$  are then chosen in a  $H^{0,s}$  neighborhood of  $u$ .

- (2) If  $u : \{t_1\} \times \mathbb{R}^d \rightarrow \mathbb{C}$ , then  $\|u(t_1)\|_{L^2} = \|v(-t_1^{-1})\|_{L^2}$ .
- (3) If  $u : [t_1, t_2] \times \mathbb{R}^d \rightarrow \mathbb{C}$  and  $(q, r)$  is admissible, then  $\|u\|_{L^q_{t \in [t_1, t_2]} L^r_x} = \|v\|_{L^q_{t \in [-t_1^{-1}, -t_2^{-1}]} L^r_x}$ .
- (4) Up to a reflection, the pseudoconformal transform is its own inverse:  $\mathcal{C}[\mathcal{C}[u]](t, x) = u(t, -x)$

These facts may be validated through explicit calculations.

## 2. GLOBAL EXISTENCE FOR INITIAL DATA IN $H^{0,s}$

The goal is to prove global existence for initial data in  $H^{0,s}$  from the assumption that there is global well-posedness in  $H^s$ .

Heuristically, initial data  $u_0 \in H^{0,s}$  at  $t_0 = 0$  can be transformed to initial data  $v_0 \in H^s$  at  $\tau_0 = -\infty$ . Under the  $H^s$  global well-posedness hypothesis,  $v$  can then be defined for all time, and  $u$  can be defined for all time by the inverse pseudoconformal transform. To make this heuristic rigorous,  $u_0$  can be evolved to  $u(T_{lwp})$  and then pseudoconformally transformed to  $v$ . Following this, it is sufficient to show that  $v(-T_{lwp}^{-1})$  is in  $H^s$  to apply the  $H^s$  global well-posedness hypothesis.

In terms of the nonlinear Schrödinger evolution map,  $S_{NLS}(t_2, t_1)$ , which was introduced earlier, the map  $u_0 \mapsto v(-T_{lwp}^{-1})$  is

$$F = \mathcal{C} \circ S_{NLS}(T_{lwp}, 0) : L^2 \rightarrow L^2.$$

Since the pseudoconformal transform commutes with the nonlinear Schrödinger evolution, this map can also be constructed in a different way, which is illustrated in Figure 1. The solutions are first allowed to evolve under  $S_{NLS}(t, 0)$  to time  $t$  (dark in left diagram), then pseudoconformally transformed from data at time  $t$  to data at transformed time  $-t^{-1}$  (dashed arrow from left diagram to right diagram), and finally allowed to evolve under  $S_{NLS}(-T_{lwp}^{-1}, -t^{-1})$  to transformed time  $-T_{lwp}^{-1}$  (dark in right diagram). This construction is represented by

$$F = F_t = S_{NLS}(-T_{lwp}^{-1}, -t^{-1}) \circ \mathcal{C} \circ S_{NLS}(t, 0) : L^2 \rightarrow L^2.$$

By the  $L^2$  local well-posedness Theorem 1.3 and the properties of the pseudoconformal transform, in a  $L^2$  neighborhood of  $u_0$ ,  $F$  is continuous with respect to the  $L^2$  norm.

To prove Proposition 1.1, it is sufficient to show that  $F$  can be restricted to  $F : H^{0,s} \rightarrow H^s$ . This is done by initially restricting to regularized data in  $H^{0,s} \cap H^1$ , showing that each of the three steps of  $F_t$  is continuous with respect to the regularized data, and then removing the regularization.  $H^{0,s} \cap H^1$  is a useful auxiliary space because it is preserved by the nonlinear Schrödinger evolution and pseudoconformally transforms to  $H^s$ .

Because the  $H^s$  local well-posedness Theorem 1.4 uses  $L^2$  Strichartz norms to control the divergence of nearby solutions, if  $u'$  and  $u''$  start near  $u$  in  $L^2$ , their separation in  $H^{0,s} \cap H^1$  can not grow by more than a constant factor. Similarly, if  $v'$  and  $v''$  start near  $v$  in  $L^2$ , then their  $H^s$  separation can not increase by more than a constant factor. Thus the divergence of the approximators, from each other, is controlled if they start in a sufficiently small  $L^2$  neighborhood of  $u_0$ .

This  $L^2$  neighborhood of  $u_0$  is illustrated by the oval in the left diagram in Figure 1. It is taken to be a ball of radius  $\delta$ , and this is the  $\delta$  which appears in the following subsections. The value of  $\delta$  is dictated by the  $H^s$  local well-posedness Theorem 1.4.

Since  $F = F_t$  is independent of  $t$ , it is possible to take the infimum in  $t$  of the  $H^s$  norm estimates for  $v'(-T_{lwp}^{-1})$  and  $v''(-T_{lwp}^{-1})$ . This eliminates the dependence on the  $H^1$  regularization and corresponds to the original heuristic of transforming  $H^{0,s}$  data at time  $t_0 = 0$  to  $H^s$  data at  $\tau_0 = -\infty$ . As a result  $F : H^{0,s} \cap H^1 \rightarrow H^s$  is continuous with respect to the  $H^{0,s}$  norm alone and extends uniquely to  $F : H^{0,s} \rightarrow H^s$  in a neighborhood of  $u_0$ . As a result, and contrary to the image in Figure 1,  $v(-T_{lwp}^{-1})$  is in  $H^s$  not merely  $L^2$ .

In Subsection 2.1, it will be shown that the nonlinear Schrödinger evolution is continuous in  $H^{0,s} \cap H^1$ . In Subsection 2.2, real interpolation will be used to show that the pseudoconformal transform takes  $H^{0,s} \cap H^1$  to  $H^s$  with a  $t$  dependent coefficient on the  $H^1$  part of the norm. In Subsection 2.3, the  $H^s$  local well-posedness theorem will be used to show that  $H^s$  data evolves continuously in  $H^s$  from transformed time  $-t^{-1}$  to transformed time  $-T_{lwp}^{-1}$ . In Subsection 2.4, the infimum in  $t$  will be taken to eliminate the  $H^1$  dependence.

2.1.  $S_{NLS}(t, 0) : H^{0,s} \cap H^1 \rightarrow H^{0,s} \cap H^1$ . By the local well-posedness theory, the nonlinear Schrödinger evolution takes  $H^1 \rightarrow H^1$ , at least up to the  $L^2$  local well-posedness time,  $T_{lwp}$ . For the linear evolution, a simple commutator calculation

shows that the  $H^1$  norm controls the growth of the  $H^{0,s}$  norm. The extra terms arising in the nonlinear evolution cancel, so that the same result holds.

**Lemma 2.1.** *For  $u$  a solution to the nonlinear Schrödinger equation (1.1) with initial data  $u_0 \in H^{0,s}$  with  $0 \leq s \leq 1$  and local existence time  $T_{lwp}$ , there is a  $\delta$  coming from the local well-posedness theory such that*

- (1) *if  $u'$  solves the nonlinear Schrödinger equation with initial data  $u'_0 \in H^{0,s} \cap H^1$  and with  $\|u_0 - u'_0\|_{L^2} < \delta$ , then  $\forall t \in [0, T_{lwp}]$*

$$\|u'(t)\|_{H^1} \leq 2\|u'_0\|_{H^1},$$

$$\|u'(t)\|_{H^{0,s}} \leq \|u'_0\|_{H^{0,s}} + 2t\|u'_0\|_{H^1}.$$

- (2) *if  $u'$  and  $u''$  solve the nonlinear Schrödinger equation with initial data  $u'_0 \in H^{0,s} \cap H^1$  and  $u''_0 \in H^{0,s} \cap H^1$  respectively and with  $\|u_0 - u'_0\|_{L^2} + \|u_0 - u''_0\|_{L^2} < \delta$ , then  $\forall t \in [0, T_{lwp}]$ ,*

$$\|u'(t) - u''(t)\|_{H^1} \leq 2\|u'_0 - u''_0\|_{H^1}$$

$$\|u'(t) - u''(t)\|_{H^{0,s}} \leq \|u'_0 - u''_0\|_{H^{0,s}} + 2t\|u'_0\|_{H^1} + 2t\|u''_0\|_{H^1}.$$

**Remark 2.2.** *The solutions  $u'$  and  $u''$  are thought of as perturbations from  $u$ . The original solution  $u$  appears in the statement of this theorem because it provides both the local well-posedness interval,  $[0, T_{lwp}]$ , on which we wish to control  $u'$  and  $u''$ , and the  $L_{tx}^{\frac{2(d+2)}{d}}$  estimates with which we can achieve this control.*

*Proof.* From the local well-posedness Theorem 1.3,  $T_{lwp}$  can be chosen small enough so that  $\|u\|_{L_{tx}^{\frac{2(d+2)}{d}}([0, T_{lwp}])}$  is less than half of  $\delta_3$  from the  $H^s$  local well-posedness

Theorem 1.4. In this case, by the  $L^2$  local well-posedness Theorem 1.3,  $\|u - u'\|_{L_{tx}^{\frac{2(d+2)}{d}}([0, T_{lwp}])} \leq 2\|u_0 - u'_0\|_{L^2} < 2\delta$  and  $\|u'\|_{L_{tx}^{\frac{2(d+2)}{d}}([0, T_{lwp}])} < 2\delta + \frac{1}{2}\delta_3$ .

The function  $u'$  can now be taken as the solution to estimate and from which  $u''$  is a perturbation. If  $2\delta + \frac{1}{2}\delta_3 \leq \delta_3$ , then the  $H^s$  local well-posedness Theorem 1.4 provides the estimates on the growth and separation in  $H^1$ .

Differentiation in time and the Cauchy-Schwartz estimate gives the growth of the weighted norms.

$$\begin{aligned} \frac{d}{dt}\|u'(t)\|_{H^{0,s}}^2 &= \frac{d}{dt}\langle u', \langle x \rangle^{2s} u' \rangle. \\ 2\|u'(t)\|_{H^{0,s}} \frac{d}{dt}\|u'(t)\|_{H^{0,s}} &= \langle i\Delta u' - i\lambda|u'|^{\frac{4}{d}}u', \langle x \rangle^{2s} u' \rangle + \langle u', \langle x \rangle^{2s} (i\Delta u' - i\lambda|u'|^{\frac{4}{d}}u') \rangle \\ &= -i\langle \Delta u', \langle x \rangle^{2s} u' \rangle + i\langle u', \langle x \rangle^{2s} \Delta u' \rangle \\ &\quad + i\lambda\langle |u'|^{\frac{4}{d}}u', \langle x \rangle^{2s} u' \rangle - i\lambda\langle u', \langle x \rangle^{2s} |u'|^{\frac{4}{d}}u' \rangle \\ &= \langle u', i[-\Delta, \langle x \rangle^{2s}]u' \rangle. \\ |2\|u'(t)\|_{H^{0,s}} \frac{d}{dt}\|u'(t)\|_{H^{0,s}}| &\leq |\langle u', -4is\langle x \rangle^{2s-2}x\Delta u' - 2is\langle x \rangle^{2s-2}((2s-2)\frac{x^2}{\langle x \rangle^2} + 1)u' \rangle| \\ &\leq C\|\langle x \rangle^{2s-1}u'\|_{L^2}\|D_x u'\|_{L^2} + C\|\langle x \rangle^{s-1}u'\|_{L^2}^2 \\ &\leq C\|u'\|_{H^{0,s}}\|D_x u'\|_{L^2} + C\|u'\|_{H^{0,s}}\|u'\|_{L^2}. \\ \frac{d}{dt}\|\langle x \rangle u'\|_{H^{0,s}} &\leq C\|D_x u'\|_{L^2} + C\|u'\|_{L^2} \\ &\leq C\|u'\|_{H^1}. \end{aligned}$$

This proves that

$$\|u'(t)\|_{H^{0,s}} \leq \|u'_0\|_{H^{0,s}} + Ct\|u'_0\|_{H^1}.$$

A similar calculation shows

$$\begin{aligned} \|u'(t) - u''(t)\|_{H^{0,s}} &\leq \|u'_0 - u''_0\|_{H^{0,s}} + \|u'(t) - u'_0\|_{H^{0,s}} + \|u''(t) - u''_0\|_{H^{0,s}} \\ &\leq \|u'_0 - u''_0\|_{H^{0,s}} + Ct(\|u'_0\|_{H^1} + \|u''_0\|_{H^1}). \end{aligned}$$

□

2.2.  $\mathcal{C} : H^{0,s} \cap H^1 \rightarrow H^s$ . In this section, it is shown that the pseudoconformal transform takes a function  $u(t) \in H^{0,s} \cap H^1$  to  $v(-t^{-1}) \in H^s$ . This is done by interpolation between  $L^2$  and  $H^{0,1} \cap H^1$  using the  $K$  method of real interpolation.

To begin, the arguments for  $L^2$  and  $H^{0,1} \cap H^1$  are presented. The  $L^2$  result is part of Theorem 1.8. The  $H^{0,1} \cap H^1$  result leads to equation (1.5), which was stated in the introduction.

**Lemma 2.3.** *If  $u : \{t\} \times \mathbb{R}^d \rightarrow \mathbb{C}$  and  $v = \mathcal{C}[u]$ , then*

$$(2.1) \quad \|v(-t^{-1})\|_{L^2} = \|u(t)\|_{L^2},$$

$$(2.2) \quad \|v(-t^{-1})\|_{H^1} \leq \|u(t)\|_{H^{0,1}} + t\|u(t)\|_{H^1}$$

where it is understood that the norm on the right is infinite (and the inequality trivial) if  $u(t)$  does not belong to the appropriate space.

*Proof.* Equation (2.1) is statement 2 of Lemma 1.8.

Inequality (2.2) follows by direct computation with a change of variables. This computation is simplified by recalling the notation  $\tau = -t^{-1}$  and  $ty = x$ .

$$\begin{aligned} \|v(-t^{-1})\|_{H^1}^2 &= \int |\nabla_y v(-t^{-1}, y)|^2 dy \\ &= \int |\nabla_y (t^{\frac{d}{2}} e^{\frac{ity^2}{4}} u(t, ty))|^2 dy \\ &= \int t^d |\frac{1}{2} itye^{\frac{ity^2}{4}} u(t, ty) + e^{\frac{ity^2}{4}} \nabla_y u(t, ty)|^2 dy \\ &\leq \int |tyu(t, ty)|^2 t^d dy + \int 2|\nabla_y u(t, ty)|^2 t^d dy \\ &\leq \int |xu(t, x)|^2 dx + 2 \int t^2 |\nabla_x u(t, x)|^2 dx. \end{aligned}$$

□

The  $K$  method of real interpolation is now summarized from [1]. The  $s$  interpolation norm of  $a \in A_0 + A_1$  is defined by the following, if this norm is finite,

$$(2.3) \quad K(\lambda, a; A_0, A_1) = \inf_{a=a_0+a_1} (\|a_0\|_{A_0}^2 + \lambda^2 \|a_1\|_{A_1}^2)^{\frac{1}{2}}$$

$$(2.4) \quad \|a\|_{s, (A_0, A_1); K}^2 = \Phi_{s,2}(K(\lambda, a; A_0, A_1)) \\ = \int \lambda^{-2s-1} K(\lambda, a; A_0, A_1)^2 d\lambda.$$

Since only the  $K$  method of interpolation will be introduced, the  $K$  index in the norm will be omitted  $\|a\|_{s, (A_0, A_1)} = \|a\|_{s, (A_0, A_1); K}$ . If  $a \in A_0 \cap A_1$ , then

$$\|a\|_{s, (A_0, A_1)} \leq \|a\|_{A_0}^{1-s} \|a\|_{A_1}^s.$$

The interpolation space  $(A_0, A_1)_s$  is defined as the set of  $a \in A_0 + A_1$  for which  $\|a\|_{s, (A_0, A_1)}$  is finite. There are some technical issues, but since only spaces  $A_0$  and  $A_1$  which are subsets of  $L^2$  will be considered,  $(A_0, A_1)_s$  will be well-defined, a Banach space, and the closure of  $A_0 \cap A_1$ .

The  $K$  method of real interpolation is an exact interpolation method of exponent  $s$  [1]: if

$$\begin{aligned} T : A_0 &\rightarrow B_0 & \|T\|_{A_0 \rightarrow B_0} &\leq M_0 \\ T : A_1 &\rightarrow B_1 & \|T\|_{A_1 \rightarrow B_1} &\leq M_1, \end{aligned}$$

then

$$(2.5) \quad \begin{aligned} T &: (A_0, A_1)_s \rightarrow (B_0, B_1)_s \\ \|T\|_{(A_0, A_1)_s \rightarrow (B_0, B_1)_s} &\leq M_0^{1-s} M_1^s. \end{aligned}$$

It is known that

$$\begin{aligned} (L^2, H^1)_s &= H^s \\ (L^2, H^{0,1})_s &= H^{0,s}. \end{aligned}$$

By interpolating the results of Lemma 2.3, it follows that

$$\mathcal{C} : (L^2, H^{0,1} \cap H^1)_s \rightarrow H^s.$$

Unfortunately, because of the inf in (2.3), it is not clear that  $\|a\|_{s, (L^2, H^{0,1} \cap H^1)} = \|a\|_{s, (L^2, H^{0,1})} + \|a\|_{s, (L^2, H^1)} = \|a\|_{H^{0,s}} + \|a\|_{H^s}$ ; although, we expect this is true. We will instead prove the simpler result that

$$\|a\|_{s, (L^2, H^{0,1} \cap H^1)} \lesssim \|a\|_{H^{0,s}} + \|a\|_{H^1}.$$

To ensure that the  $t$  dependent coefficients only appear on the  $H^1$  norm, the  $t$  dependence is kept in the interpolation calculations rather than being estimated by (2.5).

**Lemma 2.4.** *If  $u : \{t\} \times \mathbb{R}^d \rightarrow \mathbb{C}$ ,  $u(t) \in H^{0,s} \cap H^1$ , and  $v = \mathcal{C}[u]$ , then*

$$\|v(-t^{-1})\|_{H^s} \leq C_{1,s} \|u\|_{H^{0,s}} + C_{2,s} t \|u\|_{H^1}$$

where the constants  $C_{1,s}$  and  $C_{2,s}$  depend only on  $s$  and  $d$ .

*Proof.* Let  $u : \{t\} \times \mathbb{R}^d \rightarrow \mathbb{C}$  with  $u(t)$  in Schwartz class. Using the  $K$ -method, it will be shown that the  $H^s$  norm is dominated by the  $H^{0,s}$  and  $H^1$  norms. The  $K$ -method of interpolation involves taking an infimum over all possible decompositions of  $u$ . This infimum is dominated by any particular choice of decomposition. The decomposition which is optimal for balancing  $L^2$  with  $H^{0,1}$  will be used. This will give the  $H^{0,s}$  part of the estimate. There is no reduction in the regularity required for the estimate, since this decomposition ignores the  $H^1$  term.

$$\begin{aligned} \|v(-t^{-1})\|_{H^s} &= \|\mathcal{C}[u](-t^{-1})\|_{H^s} \\ &= \int \lambda^{-2s-1} K(\lambda, \mathcal{C}[u]; L^2, H^1)^2 d\lambda \end{aligned}$$

Using Lemma 2.3,  $K(\lambda, \mathcal{C}[u])$  can be estimated in terms of the  $L^2$ ,  $H^{0,1}$ , and  $H^1$  norms of  $u$ . (Note that in this proof,  $u_0$  refers to part of the interpolation

decomposition in (2.3), not the initial data.)

$$\begin{aligned} K(\lambda, \mathcal{C}[u]) &= \inf_{u=u_0+u_1} (\|\mathcal{C}[u_0]\|_{L^2}^2 + \lambda^2 \|\mathcal{C}[u_1]\|_{H^1}^2)^{\frac{1}{2}} \\ &\leq \inf_{u=u_0+u_1} (\|u_0\|_{L^2}^2 + \lambda^2 \|u_1\|_{H^{0,1}}^2 + \lambda^2 \|u_1\|_{H^1}^2)^{\frac{1}{2}}. \end{aligned}$$

In the proof that  $(L^2, H^{0,1})_s = H^{0,s}$  [1], it is shown that the optimal decomposition for  $\|u_0\|_{L^2} + \lambda^2 \|u_1\|_{H^{0,1}}^2$  is

$$\begin{aligned} u_0 &= \frac{\lambda^2(1+x^2)}{1+\lambda^2(1+x^2)}u \\ u_1 &= \frac{1}{1+\lambda^2(1+x^2)}u. \end{aligned}$$

This decomposition will be used for  $\lambda < 1$  to bound  $K(\lambda, \mathcal{C}[u])$  from above.

$$\begin{aligned} K(\lambda, \mathcal{C}[u])^2 &\leq \|u_0\|_{L^2}^2 + \lambda^2 \|u_1\|_{H^{0,1}}^2 + \lambda^2 \|u_1\|_{H^1}^2 \\ &\leq \int \left( \frac{\lambda^4(1+x^2)^2}{(1+\lambda^2(1+x^2))^2} + \frac{1}{(1+\lambda^2(1+x^2))^2} \lambda^2(1+x^2) \right) |u|^2 dx \\ &\quad + \lambda^2 t^2 \int \left| \nabla \left( \frac{1}{1+\lambda^2(1+x^2)} u \right) \right|^2 dx \\ &\leq \int \frac{\lambda^2(1+x^2)}{1+\lambda^2(1+x^2)} |u|^2 dx \\ &\quad + \lambda^2 t^2 \int \left| \nabla \left( \frac{1}{1+\lambda^2(1+x^2)} u \right) \right|^2 dx. \end{aligned}$$

This decomposition can be used to bound the  $H^s$  norm of  $v = \mathcal{C}[u]$  for  $\lambda < 1$ . The decomposition  $u_0 = u$  and  $u_1 = 0$  will be used for  $\lambda \geq 1$ .

$$\begin{aligned} \|v(-t^{-1})\|_{H^s}^2 &\leq \int_0^1 \lambda^{-2s-1} K(\lambda, \mathcal{C}[u]) d\lambda + \int_1^\infty \lambda^{-2s-1} K(\lambda, \mathcal{C}[u]) d\lambda \\ &\leq \int_0^1 \lambda^{-2s-1} \int_{\mathbb{R}^2} \frac{\lambda^2(1+x^2)}{1+\lambda^2(1+x^2)} |u|^2 dx d\lambda \\ &\quad + \int_0^1 \lambda^{-2s-1} \int \lambda^2 t^2 \left| \nabla_x \left( \frac{1}{1+\lambda^2(1+x^2)} u \right) \right|^2 dx d\lambda \\ &\quad + \int_1^\infty \lambda^{-2s-1} \|u_0\|_{L^2}^2 d\lambda \\ &\leq \int_0^1 \lambda^{-2s-1} \int_{\mathbb{R}^2} \frac{\lambda^2(1+x^2)}{1+\lambda^2(1+x^2)} |u|^2 dx d\lambda \\ &\quad + \int_0^1 \lambda^{-2s+1} t^2 \int \lambda |u|^2 + |\nabla_x u|^2 dx d\lambda \\ &\quad + C_s \|u\|_{L^2}^2. \end{aligned}$$

At this stage, the first term is evaluated by the substitution  $\lambda'^2 = (1+x^2)\lambda^2$  and Fubini's theorem. The other two pieces are estimated by direct integration and

estimated using the assumption  $t < 1$ .

$$\begin{aligned} \|v(-t^{-1})\|_{H^s}^2 &\leq \int_{\mathbb{R}^2} (1+x^2)^s |u|^2 \int_0^1 \lambda'^{-2s-1} \frac{\lambda'^2}{1+\lambda'^2} d\lambda' dx \\ &\quad + Ct^2 \|u\|_{H^1}^2 + C'(1+t^2) \|u\|_{L^2}^2 \\ &\leq C_{1,s} \|u\|_{H^{0,s}}^2 + C_{2,s} t^2 \|u\|_{H^1}^2. \end{aligned}$$

□

2.3.  $S_{NLS}(-T_{LWP}^{-1}, -t^{-1}) : H^s \rightarrow H^s$ . Since the pseudoconformal transform preserves Strichartz admissible norms,  $\|v\|_{L_{tx}^{\frac{2(d+2)}{d}}([-t^{-1}, -T_{LWP}^{-1}])}$  is controlled by  $\|u\|_{L_{tx}^{\frac{2(d+2)}{d}}([0, T_{LWP}])}$  and can be taken to be small. Since the growth of the  $H^s$  norm under the nonlinear Schrödinger evolution is controlled by  $L_{tx}^{\frac{2(d+2)}{d}}$ ,  $\|v(-T_{LWP}^{-1})\|_{H^s}$  can be controlled by  $\|v(-t^{-1})\|_{H^s}$  for any  $t \in [0, T_{LWP}]$ . The divergence of nearby solutions can be similarly controlled. From this, if  $u'_0$  and  $u''_0$  are  $H^{0,s} \cap H^1$  approximations of  $u_0 \in H^{0,s}$ , then the  $H^s$  distance between  $v'$  and  $v''$  at transformed time  $-T_{LWP}^{-1}$  is controlled by the distance between them at earlier transformed times.

**Lemma 2.5.** *For  $u$  a solution to the nonlinear Schrödinger equation (1.1) with initial data  $u_0 \in L^2$ , there is a local well-posedness time  $T_{LWP}$  and a  $\delta$  coming from the local well-posedness theory such that:*

- (1) *if  $u'$  is a solution to the nonlinear Schrödinger equation on  $[0, T_{LWP}]$  with initial data  $u'_0 \in H^{0,s} \cap H^1$ , with  $\|u_0 - u'_0\|_{L^2} < \delta$  and with a  $t \in [0, T_{LWP}]$  such that  $v'(-t^{-1}) \in H^s$ , then*

$$\|v'(-T_{LWP}^{-1})\|_{H^s} < 2\|v'(-t^{-1})\|_{H^s};$$

- (2) *if  $u'$  and  $u''$  are solutions to the nonlinear Schrödinger equation on  $[0, T_{LWP}]$  with initial data  $u'_0 \in L^2$  and  $u''_0 \in L^2$  respectively, with  $\|u_0 - u'_0\|_{L^2} < \delta$  and  $\|u_0 - u''_0\|_{L^2} < \delta$ , and with a  $t \in [0, T_{LWP}]$  such that  $v'(-t^{-1}) \in H^s$  and  $v''(-t^{-1}) \in H^s$ , then*

$$\|v'(-T_{LWP}^{-1}) - v''(-T_{LWP}^{-1})\|_{H^s} < 2\|v'(-t^{-1}) - v''(-t^{-1})\|_{H^s}.$$

**Remark 2.6.** *As in Lemma 2.1,  $u$  appears in this theorem because we want to control the size of  $L_{tx}^{\frac{2(d+2)}{d}}$  norms for the approximators,  $u'$  and  $u''$ .*

*Proof.* From the  $L^2$  local well-posedness Theorem 1.3,  $T_{LWP}$  can be chosen small enough so that  $\|u\|_{L_{tx}^{\frac{2(d+2)}{d}}([0, T_{LWP}])}$  is less than half of  $\delta_3$  from the  $H^s$  local well-posedness Theorem 1.4. In this case, by the  $L^2$  local well-posedness Theorem 1.3,  $\|u - u'\|_{L_{tx}^{\frac{2(d+2)}{d}}([0, T_{LWP}])} \leq 2\|u_0 - u'_0\|_{L^2} < 2\delta$  and  $\|u'\|_{L_{tx}^{\frac{2(d+2)}{d}}([0, T_{LWP}])} < 2\delta + \frac{1}{2}\delta_3$ .

The function  $u'$  or  $v'$  can now be taken as the solution to estimate. Since the pseudoconformal transformation preserves the Strichartz admissible norms,  $\|v'\|_{L_{tx}^{\frac{2(d+2)}{d}}([-t^{-1}, T_{LWP}^{-1}])} \leq \|v'\|_{L_{tx}^{\frac{2(d+2)}{d}}((-\infty, -T_{LWP}^{-1}])} = \|u'\|_{L_{tx}^{\frac{2(d+2)}{d}}([0, T_{LWP}])} < 2\delta + \frac{1}{2}\delta_3$ . Thus, if  $2\delta + \frac{1}{2}\delta_3 \leq \delta_3$ , then, by the  $H^s$  local well-posedness Theorem 1.4,

$$\|v'\|_{S^s([-t^{-1}, -T_{LWP}^{-1}])} < 2\|v'(-t^{-1})\|_{H^s}.$$

If  $\|u_0 - u'_0\|_{L^2} \leq \delta$  and  $\|u_0 - u''_0\|_{L^2} \leq \delta$ , then  $\|u'_0 - u''_0\|_{L^2} \leq 2\delta$ . Again, by the  $H^s$  local well-posedness Theorem 1.4, if, in addition to the previous conditions,  $2\delta \leq \delta_4$ , then

$$\|v'(-T_{lwp}^{-1}) - v''(-T_{lwp}^{-1})\|_{H^s} \leq \|v' - v''\|_{S^s([-t^{-1}, -T_{lwp}^{-1}])} \leq 2\|v'(-t^{-1}) - v''(-t^{-1})\|_{H^s}.$$

□

2.4.  $F : H^{0,s} \cap H^1 \rightarrow H^s$ . The results from Sections 2.1, 2.2, and 2.3 are now combined to show that  $F = F_t = S_{NLS}(-T_{lwp}^{-1}, -t^{-1}) \circ \mathcal{C} \circ S_{NLS}(t, 0)$  takes  $H^{0,s} \cap H^1$  to  $H^s$ . Furthermore, an explicit  $t$  dependence on the  $H^s$  norm will be found in Lemma 2.7 and then removed in Proposition 2.8 to show that  $F$  takes a  $H^{0,s}$  neighborhood of  $u_0$  into  $H^s$ .

We will first show that for  $u_0 \in H^{0,s}$ , if we restrict attention to initial data which is both in a  $H^{0,s}$  neighborhood,  $N$ , of  $u_0$  and in  $H^1$ , then  $F$  maps this initial data in  $N \cap H^1$  to  $H^s$ .

**Lemma 2.7.** *For  $u$  a solution to the nonlinear Schrödinger equation (1.1) with initial data  $u_0 \in H^{0,s}$ , then there is a local well-posedness time  $T_{lwp}$  and a  $\delta$  coming from the local well-posedness theory such that:*

- (1) *if  $u'$  is a solution to the nonlinear Schrödinger equation with initial data  $u'_0 \in H^{0,s} \cap H^1$  and  $\|u_0 - u'_0\|_{L^2} < \delta$ , then for all  $t \in (0, T_{lwp}]$  we have*

$$(2.6) \quad \|v'(-T_{lwp}^{-1})\|_{H^s} \leq C_s \|u'_0\|_{H^{0,s}} + C'_s t \|u'_0\|_{H^1};$$

- (2) *if  $u'$  and  $u''$  are solutions to the nonlinear Schrödinger equation with initial data  $u'_0 \in H^{0,s} \cap H^1$  and  $u''_0 \in H^{0,s} \cap H^1$  respectively and with  $\|u_0 - u'_0\|_{L^2} < \delta$  and  $\|u_0 - u''_0\|_{L^2} < \delta$ , then for all  $t \in (0, T_{lwp}]$  we have*

$$(2.7) \quad \|v'(-T_{lwp}^{-1}) - v''(-T_{lwp}^{-1})\|_{H^s} \leq C_s \|u'_0 - u''_0\|_{H^{0,s}} + C'_s t \|u'_0\|_{H^1} + C'_s t \|u''_0\|_{H^1}.$$

*In other words, there is an open set  $N \in H^{0,s}$  containing  $u_0$  for which*

$$F_t = S_{NLS}(-T_{lwp}^{-1}, -t^{-1}) \circ \mathcal{C} \circ S_{NLS}(t, 0) : N \cap H^1 \rightarrow H^s$$

*and  $F_t$  is continuous with respect to the  $H^{0,s} \cap H^1$  topology.*

*Proof.* Conditions on  $\delta$  and  $T_{lwp}$  will be found. To begin, assume  $T_{lwp} < 1$ .

Since  $u_0 \in H^{0,s}$  and  $u'_0$  and  $u''_0$  are  $H^{0,s} \cap H^1$  approximations, by Lemma 2.1, if  $\delta$  is less than the  $\delta$  in Lemma 2.1, then, for  $t \in [0, T_{lwp}]$ ,

$$\begin{aligned} \|u'(t)\|_{H^1} &\leq 2\|u'_0\|_{H^1}, \\ \|u'(t)\|_{H^{0,s}} &\leq \|u'_0\|_{H^{0,s}} + 2t\|u'_0\|_{H^1}, \\ \|u'(t) - u''(t)\|_{H^1} &\leq 2\|u'_0\|_{H^1} + 2\|u''_0\|_{H^1}, \\ \|u'(t) - u''(t)\|_{H^{0,s}} &\leq \|u'_0 - u''_0\|_{H^{0,s}} + 2t\|u'_0\|_{H^1} + 2t\|u''_0\|_{H^1}. \end{aligned}$$

By Lemma 2.4, the linearity of the pseudoconformal transform and the triangle inequality,

$$\begin{aligned} \|v'(-t^{-1})\|_{H^s} &\leq C\|u'_0\|_{H^{0,s}} + tC'\|u'_0\|_{H^1} \\ \|v'(-t^{-1}) - v''(-t^{-1})\|_{H^s} &\leq C\|u'_0 - u''_0\|_{H^{0,s}} + tC'\|u'_0\|_{H^1} + tC'\|u''_0\|_{H^1}. \end{aligned}$$

Since  $u_0$  is a solution with initial data in  $H^{0,s}$ ,  $u'_0$  and  $u''_0$  are  $H^{0,s} \cap H^1$  approximations, and  $v'$  and  $v''$  are in  $H^s$  at transformed time  $-t^{-1}$ , if  $T_{lwp}$  and  $\delta$  are less than the corresponding values in Lemma 2.5, then

$$\begin{aligned} \|v'(-T_{lwp}^{-1})\|_{H^s} &\leq C\|u'_0\|_{H^{0,s}} + tC'\|u'_0\|_{H^1}, \\ \|v'(-T_{lwp}^{-1}) - v''(-T_{lwp}^{-1})\|_{H^s} &\leq C\|u'_0 - u''_0\|_{H^{0,s}} + tC'\|u'_0\|_{H^1} + tC'\|u''_0\|_{H^1}. \end{aligned}$$

Since  $F_t : u_0 \mapsto v(-T_{lwp}^{-1})$  and the set  $\|u_0 - u'_0\|_{L^2} < \delta$  is open in  $H^{0,s}$ , this set is the  $N$  given in the statement of the theorem. By (2.7),  $F_t$  is continuous from  $H^{0,s} \cap H^1$  to  $H^s$ .  $\square$

The infimum in  $t$  can be taken when estimating the  $H^s$  norm of  $F_t(u_0) = v'(-T_{lwp}^{-1})$ . Since  $F = F_t : u_0 \mapsto v(-T_{lwp}^{-1})$  is independent of  $t$ , this eliminates the  $H^1$  dependence. Eliminating the  $H^1$  dependence shows that  $F$  is continuous from  $H^{0,s}$  to  $H^s$ . If  $u_0$  is approximated in  $H^{0,s}$  by a sequence of regularized initial data  $u_0^{[i]}$  in  $H^{0,s} \cap H^1$ , then the corresponding  $v^{[i]}(-T_{lwp}^{-1})$  must converge in  $H^s$  to  $v(-T_{lwp}^{-1})$ . This proves that  $v(-T_{lwp}^{-1})$  is in  $H^s$ . Under the assumption of  $H^s$  global existence,  $v$  and  $u$  can be extended globally.

**Proposition 2.8.** *Assume that the nonlinear Schrödinger equation (1.1) is globally well-posed in  $H^s$  (with the additional hypothesis that the initial data has  $L^2$  norm bounded by  $\|Q\|_{L^2}$  in the focusing case).*

*If  $u_0 \in H^{0,s}$  (and  $\|u_0\|_{L^2} < \|Q\|_{L^2}$  in the focusing case), then there is a function  $u : \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{C}$  which solves the nonlinear Schrödinger equation (1.1) for all time.*

*From the local well-posedness theory, this solution remains in  $L^2$ , at each time, with constant norm and is the unique solution in the Strichartz space  $S^0$ .*

*Proof.* For  $u_0 \in H^{0,s}$  at  $t_0 = 0$ , by Lemma 2.7, there is a solution  $u$  with local existence time  $T_{lwp}$ . If  $u'_0$  and  $u''_0$  are in  $H^{0,s} \cap H^1$  and sufficiently close to  $u_0$  in  $H^{0,s}$ , then, since the map  $F : u_0 \mapsto v(-T_{lwp}^{-1})$  is independent of  $t$ , it is possible to apply the infimum in  $t$  to (2.6) and (2.7) and obtain

$$\begin{aligned} \|v'(-T_{lwp}^{-1})\|_{H^s} &\leq C_s\|u'_0\|_{H^{0,s}}, \\ \|v'(-T_{lwp}^{-1}) - v''(-T_{lwp}^{-1})\|_{H^s} &\leq C_s\|u'_0 - u''_0\|_{H^{0,s}}. \end{aligned}$$

Since  $H^{0,s} \cap H^1$  is dense in  $H^{0,s}$ , and  $F$  is continuous with respect to the  $H^{0,s}$  norm, if a sequence  $u_0^{[i]} \in H^{0,s} \cap H^1$  is chosen to converge to  $u_0$  in  $H^{0,s}$ , the  $v^{[i]}(-T_{lwp}^{-1})$  must converge to a function  $\tilde{v} \in H^s$  and to  $v(-T_{lwp}^{-1})$  in  $L^2$ . Since  $H^s$  is dense in  $L^2$ ,  $v(-T_{lwp}^{-1}) = \tilde{v} \in H^s$ .

Since the nonlinear Schrödinger evolution and the pseudoconformal transform both preserve the  $L^2$  norm, if  $\|u_0\|_{L^2} < \|Q\|_{L^2}$ , then  $\|v(-T_{lwp}^{-1})\|_{L^2} < \|Q\|_{L^2}$ .

Therefore, in the defocusing case, from the assumption of global well-posedness in  $H^s$ ,  $v$  extends to a function  $v : \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{C}$  with  $v : \mathbb{R} \rightarrow H^s(\mathbb{R}^d)$ . For  $t > 0$ ,  $u$  can be defined by  $u = \mathcal{C}^{-1}[v]$ . By Theorem 1.8, this extension of  $u$  is a solution to the nonlinear Schrödinger equation on  $[0, \infty)$ . For  $t < 0$  all the arguments of the paper can be reproduced to define  $u$  on  $(-\infty, 0]$ . Thus,  $u$  is a solution to the nonlinear Schrödinger equation, has initial data  $u_0$ , and is defined for all  $t$ .

In the focusing case, since the nonlinear Schrödinger evolution and the pseudoconformal transform both preserve the  $L^2$  norm, if  $\|u_0\|_{L^2} < \|Q\|_{L^2}$ , then  $\|v(-T_{lwp}^{-1})\|_{L^2} <$

$\|Q\|_{L^2}$  and the same argument can be applied with the additional  $L^2$  norm hypothesis.  $\square$

**Remark 2.9.** The process of taking  $u$  on  $[0, T_{lwp}]$ , applying the pseudoconformal transform to get  $v$  on  $(-\infty, -T_{lwp}^{-1}]$ , and then extending  $v$  globally in time provides a function  $v$  which is defined for positive time. We remark that there is no clear relation between  $v$  at positive time and  $u$  at negative time. In some sense,  $v$  at positive time corresponds to the evolution of  $u$  “beyond infinite”, and, unless it is known a priori that the scattering states  $u_+$  and  $u_-$  satisfy

$$\lim_{t \rightarrow \infty} \|e^{-\frac{ix^2}{t}} e^{it\Delta} u_+ - e^{\frac{ix^2}{t}} e^{-it\Delta} u_-\|_{L^2} = 0,$$

there is no reason to believe that  $v$  at positive transformed time corresponds to  $u$  at negative time.

### 3. A SCATTERING LEMMA

**Lemma 3.1.** If  $u$  is a solution of the nonlinear Schrödinger equation (1.1) which exists globally and for which  $v$  exists globally as well, then  $\|u\|_{S^0([0, \infty))} \leq \infty$  and there exists a  $u_+$  for which

$$\lim_{t \rightarrow \infty} \|u(t) - e^{it\Delta} u_+\|_{L^2} = 0.$$

*Proof.* If  $u$  and  $v$  exist globally, then, in particular they exist for  $t \in [0, 1]$  and  $\tau \in [-1, 0]$  respectively. By the maximal time blow up Theorem 1.5,  $\|u\|_{S^0([0, 1])} < \infty$  and  $\|u\|_{S^0([1, \infty))} = \|v\|_{S^0([-1, 0])} < \infty$ . Thus  $\|u\|_{S^0([0, \infty))} < \infty$ . By Duhamel’s principle and the Strichartz estimates, this means that  $\lim_{t \rightarrow \infty} \|u(t) - e^{it\Delta} u_+\|_{L^2} = 0$ .  $\square$

**Remark 3.2.** As in Remark 2.9, it is necessary to distinguish between the forward in time extension of the pseudoconformal transform and the backwards in time extension of the pseudoconformal transform. If  $\tilde{v}$  denotes the pseudoconformal transform of  $u$  at negative time, then if  $u$  and  $\tilde{v}$  both extend globally in time, by the same argument as in this lemma,  $\|u\|_{S^0((-\infty, 0])} < \infty$  and there is a  $u_-$  for which  $\lim_{t \rightarrow -\infty} \|u - e^{-it\Delta} u_-\|_{L^2} = 0$ .

**Corollary 3.3.** Under the hypotheses of Proposition 1.1, there are functions  $u_{\pm} \in H^{0,s}$  such that

$$\lim_{t \rightarrow \pm\infty} \|e^{\mp it\Delta} u(t) - u_{\pm}\|_{H^{0,s}} = 0.$$

*Proof.* The construction in the proof of Proposition 2.8 shows that both  $u$  and  $v$  exist globally, and hence  $u$  scatters forward in time by Lemma 3.1. As noted in Remark 3.2, the same occurs backwards in time. This establishes the existence of  $u_{\pm} \in L^2$ .

We now introduce linearly advanced and retarded versions of  $u$  and  $v$ . These have two time variables, one to record the time variable associated with the nonlinear Schrödinger evolution, and one for the advancement or retardation by the linear Schrödinger evolution.

$$\begin{aligned} \phi(t, t') &= e^{i(t'-t)\Delta} u(t) \\ \psi(\tau, \tau') &= e^{i(\tau'-\tau)\Delta} v(\tau) \end{aligned}$$

The function  $\phi(t, \bullet)$  is a linear solution with initial data  $u(t)$  at time  $t' = t$ . The function  $\psi(\tau, \bullet)$  is the analogous function with initial data  $v(\tau)$  at time  $\tau' = \tau$ . Since  $v(-t^{-1})$  is the pseudoconformal transform of  $u(t)$  at time  $t$ , and the pseudoconformal transform preserves the linear Schrödinger evolution, the pseudoconformal transform of  $\phi(t, \bullet)$  with respect to the spatial variable and the second time variable is  $\psi(-t^{-1}, \bullet)$ .

Denoting a solution to the linear Schrödinger equation by  $\phi$  and its pseudoconformal transform by  $\psi$ , it is known that  $\psi(0)$  is the Fourier transform of  $\phi(0)$ . If  $u_0 \in H^{0,s}$ , by the construction in Section 2, then  $v(\tau) \in H^s$ . Since the linear Schrödinger evolution preserves the  $H^s$ , for all  $\tau'$ ,  $\psi(\tau, \tau') \in H^s$ , and, in particular  $\psi(\tau, 0) \in H^s$ . This proves that the linearly retarded version of  $u$  evolves in  $H^{0,s}$ .

$$e^{-it\Delta}u(t) = \phi(t, 0) = \mathcal{F}[\psi(\tau, 0)] \in H^{0,s}$$

Since  $e^{-it\Delta}u(t)$  evolves in  $H^{0,s}$ , it is now possible to consider scattering in  $H^{0,s}$ , despite the fact that  $u(t)$  itself does not evolve in  $H^{0,s}$ . From the  $L^2$  scattering, it follows that  $u_+ = \mathcal{F}[\psi(0, 0)] = \mathcal{F}[v(0)]$ . It now remains to show this limit holds in  $H^{0,s}$ .

$$\begin{aligned} \|e^{-it\Delta}u(t) - u_+\|_{H^{0,s}} &= \|\phi(t, 0) - \mathcal{F}[v(0)]\|_{H^{0,s}} \\ &= \|\psi(\tau, 0) - v(0)\|_{H^s} \\ &= \|e^{-i\tau\Delta}v(\tau) - v(0)\|_{H^s} \\ &= \|v(\tau) - e^{i\tau\Delta}v(0)\|_{H^s} \\ \lim_{t \rightarrow \infty} \|e^{-it\Delta}u(t) - u_+\|_{H^{0,s}} &= \lim_{\tau \rightarrow 0^+} \|v(\tau) - e^{i\tau\Delta}v(0)\|_{H^s} \rightarrow 0 \end{aligned}$$

The same argument holds as  $t \rightarrow -\infty$ , with the usual remark on the difference between the pseudoconformal transforms for positive and negative times.  $\square$

## REFERENCES

- [1] J. Bergh and J. Löfström. *Interpolation Spaces: An Introduction*. Springer-Verlag, New York, 1976.
- [2] J. Bourgain. *Global solutions of nonlinear Schrödinger equations*. AMS Colloquium Publications, 46. American Mathematical Society, Providence, R.I., 1999.
- [3] T. Cazenave, *Semilinear Schrödinger equations*, Courant Lecture Notes in Mathematics, 10. American Mathematical Society, Providence, R.I., 2003.
- [4] T. Cazenave and F. Weissler. The Cauchy problem for the critical nonlinear Schrödinger equation in  $H^s$ . *Nonlinear Anal.* 14:807-836, 1990.
- [5] M. Christ, J. Colliander, and T. Tao. Asymptotics, frequency modulation, and low regularity ill-posedness for canonical defocusing equations. *Amer. J. Math* 125:1235-1293, 2003. math.AP/0203044
- [6] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao. Almost conservation laws and global rough solutions to a nonlinear Schrödinger equation. *Math. Res. Lett.* 9:659-682, 2002. math.AP/0203218
- [7] J. Ginibre and G. Velo. The global Cauchy problem for the nonlinear Schrödinger equation revisited. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 2:309-327, 1985.
- [8] F. Linares and G. Ponce. *Introduction to Nonlinear Dispersive Equations*. *Publicações matemáticas* IMPA, 2004.
- [9] C. Sulem and P. Sulem. *Nonlinear Schroedinger Equations: Self-Focusing and Wave Collapse*. Applied Mathematical Sciences, 139. Springer, New York, 1999.
- [10] Y. Tsutsumi and K. Yajima. The asymptotic behavior of NLS equations. *Bull. Amer. Math. Soc.* 11:186-188, 1984.

- [11] M. Weinstein. Nonlinear Schrödinger equations and sharp interpolation estimates. *Comm. Math. Phys.* 87:567–576, 1983.

### Errata

In this paper, we consider the initial value problem for the  $L^2$ -critical nonlinear Schrödinger equation for  $u : \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{C}$ :

$$(3.1) \quad \begin{cases} i\partial_t u + \Delta u = \lambda |u|^{\frac{4}{d}} u \\ u(t_0, x) = u_0(x), \end{cases}$$

with initial data  $u_0$  in  $H^{0,s}$ , the weighted  $L^2$  space with norm

$$\|\psi\|_{H^{0,s}} = \left( \int_{\mathbb{R}^d} (1+x^2)^s |\psi(x)|^2 dx \right)^{1/2}.$$

Our main result was that if equation (3.1) is globally well-posed for initial data in the  $H^s$  Sobolev space, then all initial data  $u_0 \in H^{0,s}$  generates global solutions  $u \in L_t^\infty(L_x^2)$ . We also proved that such solutions scatter.

The proof presented in Section 2 approximates  $u_0 \in H^{0,s}$  by a sequence of regularised approximators in  $H^{0,s} \cap H^1$ . It has been brought to our attention that there appears to be an error in the final line of the proof of Lemma 2.1. In particular, although the growth of the  $H^{0,s}$  norms of the approximators is controlled by the  $H^1$  norm, it seems that the growth of the divergence of two approximators in  $H^{0,s}$ ,  $\|u' - u''\|_{H^{0,s}}$  is not controlled by the  $H^1$  norms of the approximators. Fortunately, the same argument works if we replace  $H^{0,s} \cap H^1$  by  $H^{0,s} \cap H^\rho$  with  $\rho = (d+1)/2$ .

In the proof of Lemma 2.1, the application of Theorem 1.4 applies equally well to the persistence of regularity in  $H^\rho$  as it did in  $H^1$  in the original proof. Thus, the same norm bound applies. The calculation of the growth of the  $H^{0,s}$  norm of  $u'$  is correct, and, since  $1 \leq (d+1)/2$ , we can trivially replace the  $H^1$  norm by the  $H^\rho$  norm. In the final calculation of Lemma 2.1, where there seems to be an error, we can still use  $\langle \cdot, \cdot \rangle$  for the  $L^2$  inner product and  $\langle x \rangle$  for  $(1+x^2)^{1/2}$  and make the estimate:

$$(3.2) \quad \begin{aligned} \frac{d}{dt} \|u'(t) - u''(t)\|_{H^{0,s}}^2 &= i \langle \Delta(u' - u''), \langle x \rangle^{2s} (u' - u'') \rangle \\ &\quad - i \langle (u' - u''), \langle x \rangle^{2s} \Delta(u' - u'') \rangle \\ &\quad - i \lambda \langle |u'|^{\frac{4}{d}} u' - |u''|^{\frac{4}{d}} u'', \langle x \rangle^{2s} (u' - u'') \rangle \\ &\quad + i \lambda \langle u' - u'', \langle x \rangle^{2s} (|u'|^{\frac{4}{d}} u' - |u''|^{\frac{4}{d}} u'') \rangle. \end{aligned}$$

The first two terms on the right can be estimated as in the estimate of the norm of  $u'$ , so that they are dominated by

$$(3.3) \quad C \|u' - u''\|_{H^{0,s}} \|u' - u''\|_{H^1}.$$

The argument which needs correcting concerns the estimate on the remaining terms. These are controlled by

$$\begin{aligned} &C |\langle |u'|^{\frac{4}{d}} u' - |u''|^{\frac{4}{d}} u'', \langle x \rangle^{2s} (u' - u'') \rangle| \\ &\leq C \|\langle x \rangle^s (u' - u'')\|_{L^2} \|\langle x \rangle^s (|u'|^{\frac{4}{d}} u' - |u''|^{\frac{4}{d}} u'')\|_{L^2} \\ &\leq C \|u' - u''\|_{H^{0,s}} \left( \|\langle x \rangle^s u'\|_{L^2} \|u'\|_{L^\infty}^{\frac{4}{d}} + \|\langle x \rangle^s u''\|_{L^2} \|u''\|_{L^\infty}^{\frac{4}{d}} \right) \\ &\leq C \|u' - u''\|_{H^{0,s}} \left( \|u'\|_{H^{0,s}} \|u'\|_{H^\rho}^{\frac{4}{d}} + \|u''\|_{H^{0,s}} \|u''\|_{H^\rho}^{\frac{4}{d}} \right). \end{aligned}$$

Here we have used that  $\rho > d/2$ , to control the  $L^\infty$  norm. By applying the earlier estimate on the growth of the  $H^\rho$  and  $H^{0,s}$  norms and restricting  $t \leq 1$ , we have the stronger bound by

$$(3.4) \quad C\|u' - u''\|_{H^{0,s}} \left( (\|u'_0\|_{H^{0,s}} + \|u'_0\|_{H^\rho}) \|u'_0\|_{H^\rho}^{\frac{4}{d}} + (\|u''\|_{H^{0,s}} + \|u''_0\|_{H^\rho}) \|u''_0\|_{H^\rho}^{\frac{4}{d}} \right).$$

Combining the estimates in (3.3) and (3.4), we have that

$$\begin{aligned} & \frac{d}{dt} \|u'(t) - u''(t)\| \\ & \leq C \left( (\|u'_0\|_{H^{0,s}} + \|u'_0\|_{H^\rho}) \left( 1 + \|u'_0\|_{H^\rho}^{\frac{4}{d}} \right) + (\|u''\|_{H^{0,s}} + \|u''_0\|_{H^\rho}) \left( \|u''_0\|_{H^\rho}^{\frac{4}{d}} + 1 \right) \right). \end{aligned}$$

Thus, the statement of Lemma 2.1 remains valid if we (i) replace the space  $H^1$  by the space  $H^\rho$ , (ii) replace the  $H^1$  norm by the  $H^\rho$  norm in estimates of the growth of Sobolev norms, and (iii) replace the  $\|u'\|_{H^1}$  norm by

$$(3.5) \quad (\|u'_0\|_{H^{0,s}} + \|u'_0\|_{H^\rho}) \left( 1 + \|u'_0\|_{H^\rho}^{\frac{4}{d}} \right)$$

and similarly for  $u''$  in estimates of the growth of the  $H^{0,s}$  norm.

In Sections 2.2 and 2.3, it is sufficient to replace  $H^1$  by  $H^\rho$ . In the proof of 2.3 and 2.4, it is sufficient to use  $H^1$  throughout and then conclude with the embedding of  $H^\rho \hookrightarrow H^1$ . In Lemma 2.5, it is sufficient to replace  $H^1$  by  $H^\rho$ , since both are sufficient to put  $v'$  and  $v''$  in  $H^s$ .

In Section 2.4, it is sufficient to make the same three changes as in Section 2.1. The replacement of the  $H^1$  norm by the term in (3.5) does not prevent the technique of taking  $t \rightarrow 0$  to leave dependence only on the  $H^{0,s}$  norm. No further use of the regularised approximators is made in the paper.

**Acknowledgements.** We would like to thank Sekin Demirbař of Bogazici University for bringing the error in the original paper to our attention.